



Did the patient have one or more of the following active diagnoses? (Validation)

SELMI = Old Myocardial Infarction ICD-9-CM code 412 = old myocardial infarction. The abstractor may determine the patient had a past AMI from clinician documentation, and presence of the 412 code is not an absolute requirement

SELPCI = PCI in past two years (Enable IHD Module) Abstractor must know approximate month and year of px ICD-9-CM Code: 00.66

SELCABG = CABG in past two years (Enable IHD Module) Abstractor must know approximate month and year of px ICD-9-CM Codes: 36.1, 36.2

VASDISC1 - VASDISC99 (PI)

Within the past two years, at any inpatient or outpatient encounter, did the patient have an active diagnosis of any of the following?
Indicate all that apply:
VASCDIS1 Coronary artery disease 414.0x, 414.2, 414.8, 414.9, 429.2
VASCDIS2 Stable angina 411.xx, 413.xx
VASCDIS3 Lower extremity arterial disease/ peripheral artery disease 440.2, 440.4
VASCDIS4 Transient cerebral ischemia 435.xx
VASCDIS5 Stroke 433.xx, 434.xx
VASCDIS6 Atheroembolism 444.xx, 445.xx
VASCDIS7 Abdominal aortic aneurysm 441.xx
VASCDIS8 Renal artery atherosclerosis 440.1
VASCDIS9 No ischemic vascular disease diagnosis

LIPSRCH = 1 (Shared)

* this question is <u>only</u> asked if patient also has diabetes * Is there documentation in the record that the patient was enrolled in a clinical trial or research protocol that precludes access to the lipid profile?

- 1. yes
- 2. no
- 95. not applicable

LDLDONE (Shared)

Was an LDL value obtained? 1. yes

- 2. no
- 2. 110

LDLDT (Shared)

Enter the date the most recent LDL value was obtained.

LDLCALC (Shared)

How was the LDL value measured?

- 1. direct
- 2. calculated
- 3. triglycerides too high to obtain valid LDL-C
- 95. not applicable

