

In Denominator Only

(Fails indicator)

# REFOPTOB (Global Measures)

Did the patient receive a referral for Outpatient Tobacco Cessation Counseling?

- The referral to outpatient tobacco cessation counseling treatment was made by the healthcare provider prior to discharge.
- Referral information was given to the patient at discharge but the appointment was not made by the provider prior to discharge.
- The referral for outpatient tobacco cessation counseling treatment was not offered because the patient's residence is not in the USA.
- 98. Patient refused the referral for outpatient tobacco cessation counseling treatment and the referral was not made.
- 99. The referral for outpatient tobacco cessation counseling treatment was not offered at discharge or unable to determine from the medical record documentation.

#### PRINCODE (Global Measures)

Enter the ICD-9-CM principal diagnosis code.

### OTHRCODE1-12 (Global Measures)

Enter the ICD-9-CM other diagnosis codes:

# TOBMEDC (Global Measures)

Was an FDA-approved tobacco cessation medication prescribed at discharge?

- A prescription for an FDA-approved tobacco cessation medication was given to the patient at discharge.
- 3. A prescription for an FDA-approved tobacco cessation medication was not offered because the patient's residence is not in the USA.
- 98. A prescription for an FDA-approved tobacco cessation medication was offered at discharge and the patient refused.
- 99. A prescription for an FDA-approved tobacco cessation medication was not offered at discharge or unable to determine from medical record documentation.

# NOTOBRXDC (Global Measures)

Is there documentation of a reason for not administering one of the FDAapproved tobacco cessation medications at discharge?

- -- Allergy to all of the FDA-approved tobacco cessation medications.
- -- Drug interaction (for all of the FDA-approved medications) with other drugs the patient is currently taking.
- -- Other reasons documented by physician/APN/PA or pharmacist.
- 1. Yes
- 2. No