

**TOBTXCOUN** (Global Measures)

Did the patient receive practical counseling that included all of the following components within the first three days of hospital admission?

- Recognizing danger situations
- Developing coping skills
- Providing basic information about quitting

1. Yes

2. No

98. Patient refused/declined practical counseling within the first three days of hospital admission

**TOBSTATUS2** (Global Measures)

What is the patient's tobacco use status documented within the first three days of admission?

1. The patient has smoked cigarettes daily on average in a volume of five or more cigarettes ( $\geq \frac{1}{4}$  pack) per day and/or cigars daily and/or pipes daily during the past 30 days
2. The patient has smoked cigarettes daily on average in a volume of four or less cigarettes ( $< \frac{1}{4}$  pack) per day and/or used smokeless tobacco and/or smoked cigarettes but not daily and/or cigars but not daily and/or pipes but not daily during the past 30 days
3. The patient has not used any forms of tobacco in the past 30 days

98. The patient refused the tobacco use screen

99. The patient was not screened for tobacco use within the first three days of admission or unable to determine the patient's tobacco use status from medical record documentation

**NOTOBMED** (Global Measures)

Is there documentation of a reason for not administering one of the FDA-approved tobacco cessation medications within the first three days of admission?

- Allergy to all of the FDA-approved tobacco cessation medications.
- Drug interaction (for all of the FDA-approved medications) with other drugs the patient is currently taking.
- Other reasons documented by physician/APN/PA or pharmacist.

1. Yes

2. No

**TOBTXMED** (Global Measures)

Did the patient receive one of the FDA-approved tobacco cessation medications within the first three days of hospital admission?

1. Yes

2. No

98. Patient refused FDA-approved tobacco cessation medications within the first three days of hospital admission