

COMFORT (STK)

When is the earliest physician, APN, or PA documentation of comfort measures only?

- Day of arrival (day 0) or day after arrival (day 1)
- 2. Two or more days after arrival (day 2 or greater)
- 3. Comfort measures only documented during hospital stay, but timing unclear
- 99. Comfort measures only was not documented by the physician/APN/PA or unable to determine

CLNTRIAL (STK)


During this hospital stay, was the patient enrolled in a clinical trial in which patients with stroke were being studied?

- 1. Yes
- 2. No

ECARINTV (STK)

Is there documentation that this admission was for the sole purpose of performance of an elective carotid intervention procedure (e.g., elective carotid endarterectomy, carotid angioplasty, carotid stenting)?

- 1. Yes
- 2. No



NOADMPRO (STK)

Is there physician/APN/PA or pharmacist documentation of a reason why VTE prophylaxis was not administered the day of or day after hospital admission?

- 1. Yes
- 2. No

95. Not applicable

ORALXAI (STK)

Is there physician/APN/PA documentation of a reason why Oral Factor Xa Inhibitor was administered for VTE prophylaxis?

- 1. Yes
- 2. No

95. Not applicable

What type of VTE prophylaxis was administered during this hospitalization? Indicate all that apply:

VTEPRO1. Low dose unfractionated heparin (LDUH), subcutaneous route only

VTEPRO2. Low molecular weight heparin (LMWH), such as enoxaparin

VTEPRO3. Intermittent pneumatic compression devices (such as SCDs)

VTEPRO4. Graduated compression stockings (such as TED hose)

VTEPRO5. Parenteral Factor Xa Inhibitor (fondaparinux)

VTEPRO6. Warfarin

VTEPRO7. Venous foot pumps (VFP)


VTEPRO8. Oral Factor Xa Inhibitor (such as rivaroxaban)

VTEPRO9. Aspirin

VTEPROA None of the above or unable to determine from medical record

documentation

- -1 Yes
- 0 No



VTESURG (VTE)

Was a surgical procedure performed using general or neuraxial anesthesia the day of or the day after hospital admission?

- 1. Yes
- 2. No

ADMPRODT (VTE)

Enter the date the VTE prophylaxis was administered after hospital admission.