

**REFOPTOB** (Global Measures)

Did the patient receive a referral for Outpatient Tobacco Cessation Counseling?

1. The referral to outpatient tobacco cessation counseling treatment was made by the healthcare provider prior to discharge.
2. Referral information was given to the patient at discharge but the appointment was not made by the provider prior to discharge.
4. The referral for outpatient tobacco cessation counseling treatment was not offered because the patient's residence is not in the USA.
98. Patient refused the referral for outpatient tobacco cessation counseling treatment and the referral was not made.
99. The referral for outpatient tobacco cessation counseling treatment was not offered at discharge or unable to determine from the medical record documentation.

PRINCODE (Global Measures)

Enter the ICD-9-CM principal diagnosis code.

OTHRCODE1-12 (Global Measures)

Enter the ICD-9-CM other diagnosis codes:

TOBMEDC (Global Measures)

Was an FDA-approved tobacco cessation medication prescribed at discharge?

1. A prescription for an FDA-approved tobacco cessation medication was given to the patient at discharge.
3. The patient's residence is not in the USA.
98. A prescription for an FDA-approved tobacco cessation medication was offered at discharge and the patient refused.
99. A prescription for an FDA-approved tobacco cessation medication was not offered at discharge or unable to determine from medical record documentation.

NOTOBRXDC (Global Measures)

Is there documentation of a reason for not administering one of the FDA-approved tobacco cessation medications at discharge?

- Allergy to all of the FDA-approved tobacco cessation medications.
 - Drug interaction (for all of the FDA-approved medications) with other drugs the patient is currently taking.
 - Other reasons documented by physician/APN/PA or pharmacist.
1. Yes
 2. No