Document Links:

Global Measures Instrument

CATNUM (rcvd on pull list)
Sample category number
70 = Global Measures

PRINCODE (Global Measures)
Enter the ICD-10-CM principal diagnosis code.

OTHRCODE1-12 (Global Measures)
Enter the ICD-10-CM other diagnosis codes:

REFOPTOB (Global Measures)
Did the patient receive a referral for Outpatient Tobacco Cessation Counseling?
1. The referral to outpatient tobacco cessation counseling treatment was made by the healthcare provider prior to discharge.
2. Referral information was given to the patient at discharge but the appointment was not made by the provider prior to discharge.
4. The referral for outpatient tobacco cessation counseling treatment was not offered because the patient's residence is not in the USA.
98. Patient refused the referral for outpatient tobacco cessation counseling treatment and the referral was not made.
99. The referral for outpatient tobacco cessation counseling treatment was not offered at discharge or unable to determine from the medical record documentation.

TOBSTATUS2 (Global Measures)
What is the patient's tobacco use status documented within the first day of admission?
1. The patient has smoked cigarettes daily on average in a volume of five or more cigarettes (>= ¼ pack) per day AND/ OR cigars daily AND/OR pipes daily during the past 30 days
2. The patient has smoked cigarettes daily on average in a volume of four or less cigarettes (< ¼ pack) per day AND/ OR used smokeless tobacco AND/OR smoked cigarettes but not daily AND/OR cigars but not daily AND/OR pipes but not daily during the past 30 days
3. The patient has not used any forms of tobacco in the past 30 days
97. The patient was not screened for tobacco use during the first day of admission because of cognitive impairment
98. The patient refused the tobacco use screen
99. The patient was not screened for tobacco use within the first day of admission or unable to determine the patient's tobacco use status from medical record documentation

TOBMEDC (Global Measures)
Was an FDA-approved tobacco cessation medication prescribed at discharge?
1. A prescription for an FDA-approved tobacco cessation medication was given to the patient at discharge.
3. The patient's residence is not in the USA.
98. A prescription for an FDA-approved tobacco cessation medication was offered at discharge and the patient refused.
99. A prescription for an FDA-approved tobacco cessation medication was not offered at discharge or unable to determine from medical record documentation.

NOTOBRXDC (Global Measures)
Is there documentation of a reason for not prescribing one of the FDA-approved tobacco cessation medications at discharge?
-- Allergy to all of the FDA-approved tobacco cessation medications.
-- Drug interaction (for all of the FDA-approved medications) with other drugs the patient is currently taking.
-- Other reasons documented by physician/APN/PA or pharmacist.
1. Yes
2. No