

Document Links:[CGPI PI Module](#)[CGPI MH Module](#)**CATNUM**

Sample category
 16. AMI - Outpatient visit
 36. SCI Dx
 48. Female, age 20-69
 50. Random Sample
 51. Random Sample MH
 54. Frail/Elderly
 60. DM Outpatient
 61. Inpatient SC
 68. Contract CBOC

REVSTAT

REVIEW STATUS (not abstracted)
 0. Abstraction has not begun
 1. Abstraction in progress
 2. Abstraction completed w/o errors
 3. TVG failure (exclusion)
 4. Record contains missing required answers
 5. Administrative exclusion from all measures

FEFLAG (rcvd on pull list)
 FE case flagged for CGPI review / scoring?
 0. No
 1. Yes

DOCHOSPCE (PI module)

Is one of the following documented in the medical record?
 -- The patient is enrolled in a VHA or community-based Hospice program
 -- The patient has a diagnosis of cancer of the liver, pancreas, or esophagus
 -- On the problem list it is documented the patient's life expectancy is less than 6 months?
 1. Yes
 2. No

DEMENTDX2 (MH)

During the past year, does the record document a diagnosis of dementia/neurocognitive disorder as evidenced by one of the following ICD-10-CM diagnosis codes:

A8100, A8101, A8109, A812, A8182, A8189, A819, F0150, F0151, F0280, F0281, F0390, F0391, F1027, F1097, F1327, F1397, F1817, F1827, F1897, F1917, F1927, F1997, G231, G300, G301, G308, G309, G3101, G3109, G3183, G903

1. Yes
 2. No

DEMSEV (MH)

Was the severity of dementia assessed during the past year using one of the following standardized tools?
 1. Clinical Dementia Rating Scale (CDR)
 2. Functional Assessment Staging Tool (FAST)
 3. Global Deterioration Scale (GDS)
 99. Severity of dementia was not assessed during the past year using one of the specified tools


COGSCOR2 (MH)

What was the outcome of the assessment of the severity of dementia assessment?

4. Score indicated mild dementia
 5. Score indicated moderate to severe dementia
 6. Score indicated no dementia
 99. No score documented in the record or unable to determine outcome

MODSEVCI (MH)

During the past year, did the clinician document in the record that the patient has moderate or severe cognitive impairment?
 1. Yes
 2. No



DEPTXYR (MH)

Within the past year, did the patient have at least one clinical encounter where depression was identified as a reason for the clinical encounter as evidenced by one of the following ICD-10-CM diagnosis codes:

F32, F320 - F325, F328, F329, F33, F330, F331, F332, F333, F334, F3340, F3341, F3342, F339, F341, F338, F0631, F0632

1. Yes
2. No

BPDXYR (MH)

Within the past year, did the patient have at least one clinical encounter where bipolar disorder was identified as a reason for the clinical encounter as evidenced by one of the following ICD-10-CM diagnosis codes:

F30, F301, F3010 - F3013, F302 - F304, F308, F309, F31, F310, F311, F3110 - F3113, F312, F313, F3130 - F3132, F314 - F316, F3160 - F3164, F317, F3170 - F3178, F318, F3181, F3189, F319

1. Yes
2. No

SCRPHQ2 (MH)

During the past year was the patient screened for depression by the PHQ-2?

1. Yes
2. No

SCRPHQI9 (MH)

During the past year was the patient screened for depression by the PHQ-2 + I9?

1. Yes
2. No
98. Patient refused depression screening

PH9SCOR (MH)

Enter the score for item 9/question #3 of the PHQ-2 + I9 screen documented in the record:

Over the last 2 weeks, how often have you been bothered by thoughts that you would be better off dead, or of hurting yourself in some way?

0. Not at all → 0
1. Several days → 1
2. More than half the days → 2
3. Nearly every day → 3
99. No answer documented

OUTCOMEI9 (MH)


Enter the interpretation of item 9/question #3 of the PHQ-2 +I9 screen as documented in the medical record.


1. Positive
2. Negative
99. No interpretation documented

CSSRS (MH)

On (computer to display **phqi9dt**), the day of the positive I 9, did the acceptable provider complete the Columbia-Suicide Severity Rating Scale (C-SSRS) Screener?

1. Yes
2. No
98. Patient refused to complete the C-SSRS screener



**CSSRSCOR3 (MH)**

Enter the score for C-SSRS Screener Question 3 documented in the record: (Time period designated, e.g. Over the past month) Have you been thinking about how you might do this?

1. Yes
2. No
95. Not applicable
99. Score not documented

CSSRSCOR4 (MH)

Enter the score for C-SSRS Screener Question 4 documented in the record: (Time period designated e.g., Over the past month) Have you had these thoughts and had some intention of acting on them?

1. Yes
2. No
99. Score not documented

CSSRSCOR5 (MH)

Enter the score for C-SSRS Screener Question 5 documented in the record: (Time period designated e.g., Over the past month) Have you started to work out or worked out the details of how to kill yourself?

1. Yes
2. No
99. Score not documented

CSSRSCOR8 (MH)

Enter the score for C-SSRS Screener Question 1 documented in the record: (Time period designated, e.g. Over the past month) Have you wished you were dead or wished you could go to sleep and not wake up?

1. Yes
2. No
99. Score not documented

OUTCOME4 (MH)

Enter the interpretation of the C-SSRS Screener as documented in the medical record.

1. Positive
2. Negative
99. No interpretation documented

VACSRA (MH)

On (computer to display **phqi9dt**), the same calendar day as the positive C-SSRS and positive I9, is there evidence of a signed VA Comprehensive Suicide Risk Assessment (CSRA) in the record?

1. Yes
2. No
98. Patient refused to complete CSRA

VACSRAACU (MH)

Enter the Clinical Impression of Acute Risk as documented in the medical record:

1. High Risk - (as evidenced by):
2. Intermediate Risk – (as evidenced by):
3. Low Risk – (as evidenced by):
99. Acute risk not documented

VACSRACHR (MH)


Enter the Clinical Impressions of Chronic Risk as documented in the medical record:

1. High Risk - (as evidenced by):
2. Intermediate Risk – (as evidenced by):
3. Low Risk – (as evidenced by):
99. Chronic risk not documented

VACSRAINT1-99 (MH)

Link to question text
(too large for this page)

Pg 8a



PTSDX (MH)

Within the past year, did the patient have at least one clinical encounter where PTSD was identified as a reason for the clinical encounter as evidenced by one of the following ICD-10-CM diagnosis codes:

F431, F4310 - F4312

1. Yes
2. No

PTSRNPC (MH)

Within the past five years, was the patient screened for PTSD using the Primary Care PTSD Screen (PC-PTSD)?

1. Yes
2. No

SCRPTSD5I9 (MH)

Within the past year, was the patient screened for PTSD using the Primary Care PTSD5 +I9?

1. Yes
2. No
98. Patient refused to complete PC-PTSD5 +I9

PTSDI9SCOR (MH)

Enter the score for item 9 question #6 of the PC-PTSD5 + I9 screen documented in the record:

Over the last 2 weeks, how often have you been bothered by thoughts that you would be better off dead, or of hurting yourself in some way?

0. Not at all → 0
1. Several days → 1
2. More than half the days → 2
3. Nearly every day → 3
99. No answer documented

PTSDI9OUT (MH)


Enter the interpretation of item 9 question #6 of the PC-PTSD5- +I9 screen as documented in the medical record.


1. Positive
2. Negative
99. No interpretation documented

PTSDCSSRS (MH)

On (computer to display pcptsd5dt), the day of the positive I 9, did the acceptable provider complete the Columbia-Suicide Severity Rating Scale (C-SSRS) Screener?

1. Yes
2. No
98. Patient refused to complete the C-SSRS screener



**PTSCSRSCOR3 (MH)**

Enter the score for C-SSRS Screener Question 3 documented in the record: (Time period designated, e.g. Over the past month) Have you been thinking about how you might do this?

- 1. Yes
- 2. No
- 95. Not applicable
- 99. Score not documented

PTSCSRSCOR4 (MH)

Enter the score for C-SSRS Screener Question 4 documented in the record: (Time period designated e.g., Over the past month) Have you had these thoughts and had some intention of acting on them?

- 1. Yes
- 2. No
- 99. Score not documented

PTSCSRSCOR5 (MH)

Enter the score for C-SSRS Screener Question 5 documented in the record: (Time period designated e.g., Over the past month) Have you started to work out or worked out the details of how to kill yourself?

- 1. Yes
- 2. No
- 99. Score not documented

PTSCSRSCOR8 (MH)

Enter the score for C-SSRS Screener Question 8 documented in the medical record: Was this within the past 3 months?

- 1. Yes
- 2. No
- 95. Not applicable
- 99. Score not documented

OUTCOME5 (MH)

Enter the interpretation of the C-SSRS Screener as documented in the medical.

- 1. Positive
- 2. Negative
- 99. No interpretation documented

PTSVACSRA (MH)

On (computer to display pcptsd5dt), the same calendar day as the positive I9, is there evidence of a signed VA Comprehensive Suicide Risk Assessment (CSRA) in the record?

- 1. Yes
- 2. No
- 98. Patient refused to complete CSRA

PTSDCSRAACU (MH)

Enter the Clinical Impression of Acute Risk as documented in the medical record:

1. High Risk - (as evidenced by):
2. Intermediate Risk - (as evidenced by):
3. Low Risk - (as evidenced by):
99. Acute risk not documented

PTSDCSRACHR (MH)


Enter the Clinical Impressions of Chronic Risk as documented in the medical record:

1. High Risk - (as evidenced by):
2. Intermediate Risk - (as evidenced by):
3. Low Risk - (as evidenced by):
99. Chronic risk not documented

PTCSRAINT1-99 (MH)

Link to question text
(too large for this page)

Pg 8b



Pg 8a

Pg 8b

(MH) Please enter the course of action documented in the record from the following list of interventions (Select all that apply). **General Strategies for Managing Risk in any setting (The provider may add additional comment/interventions as needed):**

VACSRACHR1. Initiate 9-1-1/Emergency Response Rescue
VACSRACHR2. Involuntary Hospitalization
VACSRACHR3. Voluntary Hospitalization
VACSRACHR4. Initiate one-on-one monitoring
VACSRACHR5. Initiate Health and Welfare Check
VACSRACHR6. Initiate a Hospital Transportation Plan with:
VACSRACHR7. Alert Suicide Prevention Coordinator for consideration of a Patient Record Flag Category I High Risk for Suicide
VACSRACHR8. Complete or Update Safety Plan
VACSRACHR9. Increase frequency of outpatient contacts
VACSRACHR10. Lethal Means Safety Counseling
VACSRACHR11. Obtain additional information from collateral sources
VACSRACHR12. Address barriers to treatment engagement by: [add text box]
VACSRACHR13. Address psychosocial needs by: [text box]
VACSRACHR14. Address medical conditions by: [text box]
VACSRACHR15. Connection/Referral to additional support: [text box for user to enter a name]
VACSRACHR16. Consult submitted to: [add text box for user to enter a name]
VACSRACHR17. Continue to see assigned Primary Care Provider for care
VACSRACHR18. Discussion with Veteran regarding enhancement of a sense of purpose and meaning
VACSRACHR19. Educate on smartphone VA applications (e.g. Virtual Hope Box, PTSD Coach, and Breathe2Relax)
VACSRACHR20. Education on emergency services
VACSRACHR21. Follow-up appointments: [text box]
VACSRACHR22. Initiate/refer for evidence based psychotherapy
VACSRACHR23. Involve family/support system in: [text box]
VACSRACHR24. Medication reconciliation
VACSRACHR25. Pharmacotherapy intervention to reduce suicide risk (e.g., consideration of medications shown to reduce suicide risk)
VACSRACHR26. Provide Veteran with phone number for Veteran's Crisis Line: 1-800-273-8255 (press 1)
VACSRACHR27. Reevaluate current treatment plan
VACSRACHR28. Referral to Chaplaincy/pastoral care
VACSRACHR29. Referral to peer support
VACSRACHR30. Other/Comments: [text box]
VACSRACHR99. No interventions documented by the provider

(MH) Please enter the course of action documented in the record from the following list of interventions (Select all that apply). **General Strategies for Managing Risk in any setting (The provider may add additional comment/interventions as needed):**

PTCSRACHR1. Initiate 9-1-1/Emergency Response Rescue
PTCSRACHR2. Involuntary Hospitalization
PTCSRACHR3. Voluntary Hospitalization
PTCSRACHR4. Initiate one-on-one monitoring
PTCSRACHR5. Initiate Health and Welfare Check
PTCSRACHR6. Initiate a Hospital Transportation Plan with:
PTCSRACHR7. Alert Suicide Prevention Coordinator for consideration of a Patient Record Flag Category I High Risk for Suicide
PTCSRACHR8. Complete or Update Safety Plan
PTCSRACHR9. Increase frequency of outpatient contacts
PTCSRACHR10. Lethal Means Safety Counseling
PTCSRACHR11. Obtain additional information from collateral sources
PTCSRACHR12. Address barriers to treatment engagement by: [add text box]
PTCSRACHR13. Address psychosocial needs by: [text box]
PTCSRACHR14. Address medical conditions by: [text box]
PTCSRACHR15. Connection/Referral to additional support: [text box for user to enter a name]
PTCSRACHR16. Consult submitted to: [add text box for user to enter a name]
PTCSRACHR17. Continue to see assigned Primary Care Provider for care
PTCSRACHR18. Discussion with Veteran regarding enhancement of a sense of purpose and meaning
PTCSRACHR19. Educate on smartphone VA applications (e.g. Virtual Hope Box, PTSD Coach, and Breathe2Relax)
PTCSRACHR20. Education on emergency services
PTCSRACHR21. Follow-up appointments: [text box]
PTCSRACHR22. Initiate/refer for evidence based psychotherapy
PTCSRACHR23. Involve family/support system in: [text box]
PTCSRACHR24. Medication reconciliation
PTCSRACHR25. Pharmacotherapy intervention to reduce suicide risk (e.g., consideration of medications shown to reduce suicide risk)
PTCSRACHR26. Provide Veteran with phone number for Veteran's Crisis Line: 1-800-273-8255 (press 1)
PTCSRACHR27. Reevaluate current treatment plan
PTCSRACHR28. Referral to Chaplaincy/pastoral care
PTCSRACHR29. Referral to peer support
PTCSRACHR30. Other/Comments: [text box]
PTCSRACHR99. No interventions documented by the provider