

- 1. Clinical Dementia Rating Scale (CDR)
- 2. Functional Assessment Staging Tool (FAST)
- 3. Global Deterioration Scale (GDS)

99. Severity of dementia was not assessed during the past year using one of the specified tools

MDE = Missing or Invalid Data Exclusion (data error)

Pg 2 COGSCOR2 **EXCLUDED** MIssing-COGSCOR2 (MH) What was the outcome of the assessment of the 4, 6, 99 MDE Pg 2a severity of dementia assessment? 4. Score indicated mild dementia 5. Score indicated moderate to severe dementia 6. Score indicated no dementia 99. No score documented in the record or unable MODSEVCI **EXCLUDED** Missing to determine outcome MODSEVCI (MH) During the past year, did the clinician document Pg 2b in the record that the patient has moderate or severe cognitive impairment? 1. Yes 2. No **STDYBEG** Missing-MDE MDE Missing Valid Date **MDE** > 1 year Calculate Valid <= 1 year RECDEPDT **DEPTXYR** Missing-STDYBEG - RECDEPDT Date 2 **EXCLUDED** Calculate Valid <= 1 year RECBPDT **BPDXYR** STDYBEG - RECBPDT Date > 1 year ż MIssing MDE Pg 3

DEPTXYR (MH)

Within the past year, did the patient have at least one clinical encounter where depression was identified as a reason for the clinical encounter as evidenced by one of the following ICD-10-CM diagnosis codes:

F01.51, F32.0 - F32.5, F32.81, F32.89, F32.9, F33.0 - F33.3, F33.42, F33.9, F34.1, F43.21, F43.23

1. Yes

2. No

RECDEPDT (MH)

Enter the date within the past year of the <u>most recent</u> clinical encounter where depression was identified as a reason for the clinical encounter.

BPDXYR (MH)

Within the past year, did the patient have at least one clinical encounter where bipolar disorder was identified as a reason for the clinical encounter as evidenced by one of the following ICD-10-CM diagnosis codes:

F30.10 – F30.13, F30.2 – F30.4, F30.8, F30.9, F31.0, F31.10 – F31.13, F31.2, F31.30 – F31.32, F31.4, F31.5, F31.60 – F31.64, F31.70 – F31.78, F31.81, F31.89, F31.9

1. Yes

2. No

RECBPDT (MH)

Enter the date within the past year of the <u>most recent</u> clinical encounter where bipolar disorder was identified as a reason for the clinical encounter.

SCRPHQI9 (MH)

During the past year was the patient screened for depression by the PHQ-2 + I9?

- 1. Yes
- 2. No
- 98. Patient refused depression screening

PHQI9DT (MH)

Enter the date within the past year of the <u>most</u> recent screening for depression by the PHQ-2 +I9

SCRPHQ2 (MH)

During the past year was the patient screened for depression by the PHQ-2?

- 1. Yes
- 2. No

98. Patient refused depression screening by the PHQ-2

PHQ2DT (MH)

Enter the date of the <u>most recent</u> screening for depression by the PHQ-2.

PHQ1SCOR (MH)

Enter the score for PHQ-2 Question 1 documented in the record:

Over the past 2 weeks, have you been bothered by little interest or pleasure in doing things?

- 0. Not at all \rightarrow 0
- 1. Several days → 1
- 2. More than half the days \rightarrow 2
- 3. Nearly every day \rightarrow 3
- 99. No answer documented

PHQ2SCOR (MH)

Enter the score for PHQ-2 Question 2 documented in the record:

Over the past 2 weeks, have you been bothered by feeling down, depressed, or hopeless?

- 0. Not at all \rightarrow 0
- 1. Several days → 1
- 2. More than half the days \rightarrow 2
- 3. Nearly every day \rightarrow 3
- 99. No answer documented

PHQTOTAL (MH)

Enter the <u>total score</u> for the PHQ-2 questions documented in the medical record.

OUTCOME3 (MH)

Enter the interpretation of the depression screen as documented in the medical record.

- 1. Positive
- 2. Negative
- 99. No interpretation documented

