

Document Links:[HBPC Instrument](#)**COHORT**

69 – Home Based Primary Care

REVSTAT

REVIEW STATUS (not abstracted)

- 0. Abstraction has not begun
- 1. Abstraction in progress
- 2. Abstraction completed w/o errors
- 3. TVG failure (exclusion)
- 4. Record contains missing data
- 5. Administrative exclusion from all measures

HOSPICE

During the past year is there documentation in the medical record the patient is enrolled in a VHA or community-based hospice program?

- 1. Yes
- 2. No

HBPCDT

Enter the date of the most recent home care encounter for this patient, occurring within the study interval.


DEMENTDX2


During the past year, does the record document a diagnosis of dementia/neurocognitive disorder as evidenced by one of the following ICD-10-CM diagnosis codes:

A81.00, A81.01, A81.09, A81.2, A81.82, A81.89, A81.9, F01.50, F01.51, F02.80, F02.81, F03.90, F03.91, F10.27, F10.97, F13.27, F13.97, F18.17, F18.27, F18.97, F19.17, F19.27, F19.97, G23.1, G30.0, G30.1, G30.8, G30.9, G31.01, G31.09, G31.83, G90.3

- 1. Yes
- 2. No

**MDE = Missing or
Invalid Data Exclusion
(data error)**



**PERMCI**

During the past year, did a physician/APN/PA or psychologist document that the patient has probable permanent cognitive impairment using a Clinical Reminder?

1. Yes
2. No

MODSEVCI

During the past year, did the clinician document in the record that the patient has moderate or severe cognitive impairment?

1. Yes
2. No

DEMSEV

Was the severity of dementia assessed during the past year using one of the following standardized tools?

1. Clinical Dementia Rating Scale (CDR)
2. Functional Assessment Staging Tool (FAST)
3. Global Deterioration Scale (GDS)
99. Severity of dementia was not assessed during the past year using one of the specified tools

COGSCOR2


What was the outcome of the assessment of the severity of dementia assessment?

4. Score indicated mild dementia
5. Score indicated moderate to severe dementia
6. Score indicated no dementia
99. No score documented in the record or unable to determine outcome

INCSEVCI

During the timeframe from (computer display demsevd + 1 day to stdyend), did a physician/APN/PA or psychologist document in the record that the patient has moderate or severe cognitive impairment?

1. Yes
2. No

**VACSRE**

On (if vacssrs = 1, computer to display vacssrsdt; else display, During the past year), is there evidence of a signed Comprehensive Suicide Risk Evaluation (CSRE) in the record?

- 1. Yes
- 2. No
- 98. Patient refused to complete CSRE

CSREACU

Enter the Clinical Impression of Acute Risk as documented in the medical record:

- 1. High Risk - (as evidenced by):
- 2. Intermediate Risk – (as evidenced by):
- 3. Low Risk – (as evidenced by):
- 99. Acute risk not documented

CSRECHR

Enter the Clinical Impression of Chronic Risk as documented in the medical record:

- 1. High Risk - (as evidenced by):
- 2. Intermediate Risk – (as evidenced by):
- 3. Low Risk – (as evidenced by):
- 99. Chronic risk not documented

STDYBEG (Rcvd on Pull List)
Study Interval begin date


VACSSRS

During the past year, did an acceptable provider complete the Columbia-Suicide Severity Rating Scale (C-SSRS) Screener?

- 1. Yes
- 2. No
- 98. Patient refused to complete the C-SSRS Screener

VACSSRSDT

Enter the most recent date the C-SSRS Screener was completed.



1. Positive
2. Negative
99. No interpretation documented



CSREINT1. Alert Suicide Prevention Coordinator for consideration of a Patient Record Flag Category I High Risk for Suicide

CSREINT2. Complete or Update Veteran's Safety Plan

CSREINT3. Increased frequency of Suicide Risk Screening
[text box]

CSREINT4. Provide Lethal Means Safety Counseling (e.g., provision of gun locks)

CSREINT5. Obtain additional information from collateral sources
[Optional: comment]

CSREINT6. For prescribers only: Review of prescribed medications for risk for self-harm and/or new pharmacotherapy intervention to reduce suicide risk
[Optional: comment]

CSREINT7. Address barriers to treatment engagement by:
[text box]

CSREINT8. Address psychosocial needs by: [text box]

CSREINT9. Address medical conditions by: [text box]

CSREINT10. Consult/Referral to additional services and support:
[text box for options]

CSREINT11. Referral to evidence based psychotherapy

CSREINT12. Referral to psychiatry/medication assessment or management

CSREINT13. Referral to Chaplaincy/pastoral care

5 of 5