

- 0. Abstraction has not begun
- 1. Abstraction in progress
- 2. Abstraction completed w/o errors
- 3. Cohort validation failure (exclusion)
- 4. Record contains missing data
- 5. Administrative exclusion from all measures

SEENYR2 (Validation)

During the timeframe from (computer display stdybeg – 1 year to stdyend),was the Veteran seen by a physician, NP, PA, Psychologist, or Clinical Nurse Specialist in one of the "Nexus clinics"?

- 1. Yes
- 2. No

Enter the date of the most recent visit to a Nexus clinic during which the patient was seen by a physician, NP, PA, Psychologist, or Clinical Nurse Specialist.


VALNEXUS (Validation)

On (computer to display pnexusdt), is there documentation the patient was seen by a physician, NP, PA, Psychologist, or Clinical Nurse Specialist in one of the "Nexus clinics"?

- 1. Yes
- 2. No

PNEXUSDT (Validation)
Computer will prefill the date
of the most recent visit to a
Nexus clinic during which the
patient was seen by a
physician, NP, PA,
Psychologist, or Clinical Nurse
Specialist.

MDE = Missing or Invalid Data Exclusion (data error)



PERMCI (MH)

During the past year, did a physician/APN/PA or psychologist document that the patient has probable permanent cognitive impairment using a Clinical Reminder?

- 1. Yes
- 2. No

HOSPICE (Validation)

During the past year is there documentation in the medical record the patient is enrolled in a VHA or community-based hospice program?


- 1. Yes
- 2. No

DEMENTDX2 (MH)

During the past year, does the record document a diagnosis of dementia/neurocognitive disorder as evidenced by one of the following ICD-10-CM diagnosis codes:

A81.00, A81.01, A81.09, A81.2, A81.82, A81.89, A81.9, F01.50, F01.51, F02.80, F02.81, F03.90, F03.91, F10.27, F10.97, F13.27, F13.97, F18.17, F18.27, F18.97, F19.17, F19.27, F19.97, G23.1, G30.0, G30.1, G30.8, G30.9, G31.01, G31.09, G31.83, G90.3

- 1. Yes
- 2. No



DEMSEV (MH)

Was the severity of dementia assessed during the past year using one of the following standardized tools?

- 1. Clinical Dementia Rating Scale (CDR)
- 2. Functional Assessment Staging Tool (FAST)
- 3. Global Deterioration Scale (GDS)
- 99. Severity of dementia was not assessed during the past year using one of the specified tools

COGSCOR2 (MH)

What was the outcome of the assessment of the severity of dementia assessment?

- 4. Score indicated mild dementia
- Score indicated moderate to severe dementia
- 6. Score indicated no dementia
- 99. No score documented in the record or unable to determine outcome

During the timeframe from (computer display demsevdt + 1 day to stdyend), did a physician/APN/PA or psychologist document in the record that the patient has moderate or severe cognitive impairment?

- 1. Yes
- 2. No

MODSEVCI (MH)


During the past year, did the clinician document in the record that the patient has moderate or severe cognitive impairment?

- 1. Yes
- 2. No

VACSSRS (MH)

During the past year, did an acceptable provider complete the Columbia-Suicide Severity Rating Scale (C-SSRS) Screener?

- 1. Yes
- 2. No
- 98. Patient refused to complete the C-SSRS Screener



CSSRSOUT (MH)


Enter the interpretation of the C-SSRS Screener as documented in the medical record.

- 1. Positive
- 2. Negative
- 99. No interpretation documented

ADMINPT (MH)

On (computer to display vacssrsdt), the same calendar day as the positive C-SSRS, is there evidence the patient was admitted to inpatient or residential treatment for mental health care?

- 1. Yes
- 2. No



of a Patient Record Flag Category I High Risk for Suicide

CSREINT2. Complete or Update Veteran's Safety Plan

CSREINT3. Increased frequency of Suicide Risk Screening [text box]

CSREINT4. Provide Lethal Means Safety Counseling (e.g., provision of gun locks)

CSREINT5. Obtain additional information from collateral sources [Optional: comment]

CSREINT6. For prescribers only: Review of prescribed medications for risk for self-harm and/or new pharmacotherapy intervention to reduce suicide risk (Optional: comment)

CSREINT7. Address barriers to treatment engagement by: [text box]

CSREINT8. Address psychosocial needs by: [text box]

CSREINT9. Address medical conditions by: [text box]

CSREINT10. Consult/Referral to additional services and support: [text box for options]

CSREINT11. Referral to evidence based psychotherapy

CSREINT12. Referral to psychiatry/medication assessment or management

CSREINT13. Referral to Chaplaincy/pastoral care

CSREINT15. Referral for PRRC and/or ICMHR services

CSREINT16. Referral for residential mental health services

CSREINT17. Other Consult submitted to: [text box for user to enter a name]

CSREINT18. Discussion with Veteran to continue to see assigned Primary Care Provider for medical care

CSREINT19. Discussion with Veteran regarding enhancement of a sense of purpose and meaning

CSREINT20. Educate Veteran on smartphone VA applications (e.g. Virtual Hope Box, PTSD Coach)

CSREINT21. Conduct medication reconciliation

CSREINT22. Involve family/support system in Veteran's care

CSREINT23. Provide Opioid Overdose Education and Naloxone Distribution (OEND)

CSREINT24. Provide resources/contacts for benefits information CSREINT25. Provide Veteran with phone number for Veteran's

Crisis Line: 1-800-273-8255 (press 1)

CSREINT26. Other/Comments: [text box]

CSREINT27. Obtain consultation from Suicide Risk Management Consultation Program on ways to address Veteran's risk by sending a request for consultation by email to: Email (Left Click and Allow)

CSREINT99. No interventions documented by the provider