

Meets

denominator of Sep1b

Yes

Meets

numerator of


Sep1b

Yes ↓ Pg 2

No

EXCLUDED

No-



HYPOTNS

During the time frame from (computer to display sepresdt/seprestm - 6 hours) to (computer to display sepresdt/seprestm + 6 hours) is there documentation initial hypotension was present? Criteria for determining initial hypotension:

- Two hypotensive blood pressure readings at different times within specified timeframe o systolic blood pressures <90, or

 - o mean arterial pressures (MAP), <65 or o a decrease in systolic BP by >40 mm/Hg
- 1. Yes
- 2. No or unable to determine

HYPOTNSDT / HYPOTNSTM

Enter the date and time on which initial hypotension was present during the time frame from(computer to display sepresdt/seprestm - 6 hours to sepresdt/ seprestm + 6 hours).

CRYSTL2

During the timeframe from [(If hypotns = 1, and sepshk = 2, computer to display hypotnsdt/hypotnstm - 6 hours) to (hypotnsdt/hypotnstm + 3 hours) OR (If hypotns = 2 and sepshk = 1 computer to display sepshkdt/sepshktm - 6 hours) to (sepshkdt/sepshktm + 3 hours) OR (if hypotns = 1 and sepshk = 1, computer to display earliest of hypotnsdt/hypotnstm - 6 hrs) to (hypotnsdt/hypotnstm + 3 hours or sepshkdt/sepshktm - 6 hours to sepshkdt/ sepshktm + 3 hours) were crystalloid fluids initiated?

- 1. Yes
- 2. No

4. There is documentation the patient has an implanted Ventricular Assist Device (VAD)

- 97. Physician/APN/PA or nursing documentation no fluids were ordered because the patient was not volume or fluid responsive by clinical evidence.
- 98. The patient or authorized patient advocate refused IV fluids.

TARGVOL

Is there documentation the target ordered volume of crystalloid fluids (computer to display crystlvol) initiated on (computer to display crystldt at crystltm) was completely infused?

- 1. Yes
- 2. No

CRYSTLDT / CRYSTLTM

Enter the earliest date and time on which crystalloid fluids were initiated.

PERSHYPO

During the time frame from (computer display crystlend) to (computer to display crystlend + 1 hour) is there physician/APN/PA documentation that persistent hypotension or new onset of hypotension was present?

Criteria for determining persistent or new onset of hypotension:

• Two hypotensive blood pressure readings at different times within specified timeframe


- o systolic blood pressures <90, or
- o mean arterial pressures (MAP), <65 or
- o a decrease in systolic BP by >40 mm/Hg

1. Yes

- 2. No or Unable to determine
- 3. No, the patient was not assessed for persistent hypotension or new onset of hypotension within one hour after the conclusion of crystalloid fluid administration at

the target ordered volume.

4. Not applicable - Crystalloid fluids were administered but at a volume less than the target ordered volume



SEPSHK

Did a physician/APN/PA document presence of septic shock, severe sepsis with shock, or sepsis with shock?

1. Yes

2. No or unable to determine

SEPSHKDT / SEPSHKTM

Enter the earliest date and time a physician/ APN/PA documented the presence of septic shock OR the earliest date on which the final criterion was met to establish the presence of septic shock.

CNTRASEPSHK

During the timeframe from (computer to display sepshktm - 6 hours) to (computer to display sepshktm + 6 hours) is there physician/ APN/PA or nursing documentation that the patient or authorized patient advocate refused either a blood draw, IV fluid administration, or vasopressor administration?

- 1. Yes
- 2. No or unable to determine

CMOPLC2

During the timeframe from (computer to display sepshktm - 6 hours) to (computer to display sepshktm + 6 hours) is there physician/APN/PA documentation of comfort measures only or palliative care (see Inclusion terms below)?

Inclusion (Or synonymous terms)	
Hospice care	
Organ harvest	
Palliative Care	
Organ harvest	
Terminal care	
Terminal extubation	
Withdraw care	
Withhold care	

