

REVSTAT
 REVIEW STATUS (calculated)
 0. Abstraction has not begun
 1. Abstraction in progress
 2. Abstraction completed w/o errors
 3. TVG failure (exclusion)
 4. Record contains missing required answers (error record)
 5. Administrative Exclusion


DCDT (Global Measures)
 Discharge date (rcvd on pull list and may not be modified)

ADMDT (Global Measures)
 Admission date:

BIRTHDT (Global Measures)
 Patient date of birth

AGE (Calculated)
 ADMDT - BIRTHDT

COGIMP (Global Measures)
 Is there documentation in the medical record that indicates the patient is cognitively impaired?
 1. Yes
 2. No




TOBUSE (Global Measures)
 Is there documentation the patient used any form of tobacco during the past 30 days?
 1. Yes, patient has used tobacco during the past 30 days
 2. No, patient has not used tobacco during past 30 days
 98. Patient refused tobacco use screen
 99. The patient was not screened for tobacco use during this hospitalization or unable to 1, determine the patient's tobacco use status from medical record documentation.

PRINCIDE (Global Measures)
 Enter the ICD-9-CM principal diagnosis code.

OTHRCODE1 – OTHRCODE12 (Global Measures)
 Enter the ICD-9-CM other diagnosis codes:

TOBSTATUS (Global Measures)
 What is the patient's tobacco use status?
 1. Has smoked cigarettes on average in a volume of more than five cigarettes per day during the past 30 days.
 2. Has smoked cigarettes on average in a volume of five or less cigarettes per day during the past 30 days.
 3. Smokes cigarettes but does not smoke daily.
 4. Has used smokeless tobacco products in the past 30 days.
 5. Has smoked a pipe or cigar daily in the past 30 days.
 9. Has smoked cigarettes on average in a volume of more than five cigarettes per day during the past 30 days and has used smokeless tobacco products in the past 30 days.
 10. Has smoked cigarettes on average in a volume of more than five cigarettes per day during the past 30 days, has used smokeless tobacco products in the past 30 days, and has smoked a pipe or cigars daily in the past 30 days
 11. Has smoked cigarettes on average in a volume of five or less cigarettes per day during the past 30 days and has used smokeless tobacco products in the past 30 days.
 12. Has smoked cigarettes on average in a volume of five or less cigarettes per day during the past 30 days and has smoked a pipe or cigars daily in the past 30 days, and has used smokeless tobacco products in the past 30 days.
 13. Has smoked cigarettes on average in a volume of five or less cigarettes per day during the past 30 days and has smoked a pipe or cigars daily in the past 30 days.
 14. Has smoked cigarettes on average in a volume of more than five cigarettes per day during the past 30 days and has smoked a pipe or cigars daily in the past 30 days.

TOBTXCOUN (Global Measures)
 Did the patient receive practical counseling that included all of the following components prior to discharge?
 -- Recognizing danger situations
 -- Developing coping skills
 -- Providing basic information about quitting
 1. Yes
 2. No
 98. Patient refused/declined practical counseling



TOBTXMED (Global Measures)
 Did the patient receive one of the FDA-approved tobacco cessation medications during the hospital stay?
 1. Yes
 2. No
 98. Patient refused FDA-approved tobacco cessation medications during the hospital stay

NOTOBMED (Global Measures)
 Is there documentation of a reason for not administering one of the FDA-approved tobacco cessation medications during the hospital stay?
 ● Allergy to all of the FDA-approved tobacco cessation medications.
 ● Drug interaction (for all of the FDA-approved medications) with other drugs the patient is currently taking.
 ● Other reasons documented by physician/APN/PA or pharmacist.
 1. Yes
 2. No