CATNUM
Sample category
16. AMI - Outpatient visit
36. SCI Dx
48. Female, age 20-69
50. Random Sample
51. Random Sample MH
54. Frail/Elderly
60. DM Outpatient
61. Inpatient SC
68. Contract CBOC

REVSTAT
REVIEW STATUS (not abstracted)
0. Abstraction has not begun
1. Abstraction in progress
2. Abstraction completed w/o errors
3. TVG failure (exclusion)
4. Record contains missing required answers (error record)
5. Administrative exclusion from all measures

FEFLAG (rcvd on pull list)
FE case flagged for CGPI review / scoring?
0. No
1. Yes

DEMENTDX2 (MH)
During the past year, does the record document a diagnosis of dementia/neurocognitive disorder as evidenced by one of the following ICD-9-CM codes?
(046.1, 046.11, 046.19, 046.3, 290.0, 290.10, 290.11, 290.12, 290.13, 290.20, 290.21, 290.3, 290.40, 290.41, 290.42, 290.43, 291.2, 292.82, 294.10, 294.11, 294.8, 331.0, 331.11, 331.19, 331.2, 331.7, 331.82, 331.89, 331.9, 333.0 or 333.4)
1. Yes
2. No

DEMSEV (MH)
Was the severity of dementia assessed during the past year using one of the following standardized tools?
1. Clinical Dementia Rating Scale (CDR)
2. Functional Assessment Staging Tool (FAST)
3. Global Deterioration Scale (GDS)
99. Severity of dementia was not assessed during the past year using one of the specified tools

COGSCOR2 (MH)
What was the outcome of the assessment of the severity of dementia assessment?
4. Score indicated mild dementia
5. Score indicated moderate to severe dementia
6. Score indicated no dementia
99. No score documented in the record or unable to determine outcome

MODSEVCI (MH)
During the past year, did the clinician document in the record that the patient has moderate or severe cognitive impairment?
1. Yes
2. No

DEPTXYR (MH)
Within the past year, did the patient have at least one clinical encounter where depression was identified as a reason for the visit as evidenced by one of the following ICD-9-CM codes:
296.2-296.3, 298.0, 300.4, 301.12, 309.0, 309.1, 309.28, or 311.
1. Yes
2. No

BPDXYR (MH)
Within the past year, did the patient have at least one clinical encounter where bipolar disorder was identified as a reason for the clinical encounter as evidenced by one of the following ICD-9-CM codes:
296.5x, 296.6x, 296.7, and 296.8x
1. Yes
2. No

MDE = Missing or Invalid Data Exclusion (data error)

Pg 2

EXCLUDED ◄— 99/99/9999 — PHQ2DT — MIssing ► MDE

Valid Date

3 — SCRNPHQ — MIssing ►

2 or 4

99 — PH1SCOR — Missing

0, 1, 2, 3

PH1SCOR — 3

MDE

MIssing

0, 1, 2

PH2SCOR ◄— 99 or Missing — PH2SCOR — 3 ►

99

0, 1, 2

Calculate sum of PH1SCOR + PH2SCOR — >= 3 ►

< 3

MDE ◄— < 0 or MIssing — PHQTOTAL — >=3 ►

(>= 0 and < 3) or z

Pg 3 ◄— 2, 99 — OUTCOME3 — 1 ► Pg 3b

MIssing

MDE

PHQ2DT   (MH)
Enter the date within the past year of the most recent screening for depression by the PHQ-2 or PHQ-9.

SCRNPHQ (MH)
On the date of the most recent screening for depression, was the patient screened by the PHQ-2 or the PHQ-9?
2. Screened by PHQ-2
3. Screened by PHQ-9
4. Screened by PHQ-2 AND PHQ-9 on the same date

PH1SCOR (MH)
Enter the score for PHQ-2 Question 1 documented in the record:
Over the past 2 weeks, have you been bothered by little interest or pleasure in doing things?
0. Not at all --> 0
1. Several days --> 1
2. More than half the days --> 2
3. Nearly every day --> 3
99. No answer documented

PH2SCOR   (MH)
Enter the score for PHQ-2 Question 2 documented in the record:
Over the past 2 weeks, have you been bothered by feeling down, depressed, or hopeless?
0. Not at all --> 0
1. Several days --> 1
2. More than half the days --> 2
3. Nearly every day --> 399. No answer documented

PHQTOTAL   (MH)
Enter the total score for the **PHQ-2** documented in the medical record.

OUTCOME3   (MH)
What was the outcome of the PHQ-2 documented in the record?
1. Outcome positive (suggestive of depression)
2. Outcome negative (no indication of depression)
99. Outcome not documented

Pg 3

EXCLUDED ◄— 2 — SCRNPHQ

3 or 4

>= 10 and <= 27 — PH9TOTAL — < 0, > 27 or MIssing → MDE

(>= 0 and < 10) or zz

1, 2, 3 — PH9SCOR — MIssing →

0, 95, 99

1, 5, 6, 7 — PHQ9OUT — MIssing →

Pg 3b

3, 4, 95, 99

EXCLUDED

MDE ◄— MIssing — DEPRISK — 2 →

In Denominator Only
(**Fails** indicator)

1

Calculate
DEPRSKDT - PHQ2DT — < 0 or > 1

>= 0 and <= 1

In Denominator and
Numerator
(**Passes** indicator)

PHQ9TOTAL (MH)
Enter the total score of the PHQ-9 documented in the record.

PH9SCOR (MH)
Enter the score for PHQ-9 Question 9 documented in the record:
Over the last 2 weeks, how often have you been bothered by thoughts that you would be better off dead, or of hurting yourself in some way?
0.   Not at all --> 0
1.   Several days --> 1
2.   More than half the days --> 2
3.   Nearly every day --> 3
95. Not applicable
99. No answer documented

PHQ9OUT (MH)
Was the outcome of the PHQ-9 documented in the medical record?
1.   Outcome positive
3.   Score suggestive of no depression
4.   Score suggestive of mild depression
5.   Score suggestive of moderate depression
6.   Score suggestive of moderately severe depression
7.   Score suggestive of severe depression
95. Not applicable
99. No documentation of outcome

DEPRISK (MH)
Following the positive PHQ-2 or PHQ-9 depression screen or affirmative answer to PHQ-9 question 9, did the provider document a suicide ideation/behavior evaluation?
1. Yes
2. No

DEPRSKDT (MH)
Enter the date of the suicide ideation/behavior evaluation.